

# Summary Document for Interim Clinical Considerations

for Use of COVID-19 Vaccines Currently Authorized or Approved in the United States



## COVID-19 vaccine products currently approved or authorized in the United States

Pfizer-BioNTech							
Age indication	Vaccine vial cap color	Label border color	Dilution required	Primary series		Booster doses	
				Dose	Injection volume	Dose	Injection volume
6 months–4 years	Maroon	Maroon	Yes	3 µg	0.2 mL	NA	NA
5–11 years	Orange	Orange	Yes	10 µg	0.2 mL	10 µg	0.2 mL
12 years and older	Purple	Purple	Yes	30 µg	0.3 mL	30 µg	0.3 mL
12 years and older	Gray	Gray	No	30 µg	0.3 mL	30 µg	0.3 mL
Moderna							
Age indication	Vaccine vial cap color	Label border color	Dilution required	Primary series		Booster doses	
				Dose	Injection volume	Dose	Injection volume
6 months–5 years	Dark blue	Magenta	No	25 µg	0.25 mL	NA	NA
6–11 years	Dark blue	Purple	No	50 µg	0.5 mL	NA	NA
12–17 years	Red	Light blue	No	100 µg	0.5 mL	NA	NA
18 years and older	Red	Light blue	No	100 µg	0.5 mL	50 µg	0.25 mL
18 years and older	Dark blue	Purple	No	NA	NA	50 µg	0.5 mL
Janssen							
Janssen COVID-19 Vaccine is authorized for adults ages 18 years and older in certain limited situations due to safety considerations. For guidance on respective record review, scheduling and administration of Janssen vaccine see Interim Clinical Considerations for Use of COVID-19 Vaccines: Appendix A ( <a href="https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us-appendix.html#appendix-a">https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us-appendix.html#appendix-a</a> )							
Age indication	Vaccine vial cap color	Label border color	Dilution required	Primary series		Booster doses	
				Dose	Injection volume	Dose	Injection volume
18 years and older	Blue	No Color	No	5×10 <sup>10</sup> viral particles	0.5 mL	5×10 <sup>10</sup> viral particles	0.5 mL
Novavax							
Age indication	Vaccine vial cap color	Label border color	Dilution required	Primary series		Booster doses	
				Dose	Injection volume	Dose	Injection volume
12 years and older	Royal blue	No Color	No	5 µg rS and 50 µg of Matrix-M™ adjuvant	0.5 mL	N/A	N/A

# Summary Document for Interim Clinical Considerations

for Use of COVID-19 Vaccines Currently Authorized or Approved in the United States



## All currently authorized or approved COVID-19 vaccines

<b>COVID-19 vaccination schedule</b>	<ul style="list-style-type: none"> <li>See the Interim COVID-19 Immunization Schedule for Ages 6 Months or Older at <a href="https://www.cdc.gov/vaccines/covid-19/downloads/COVID-19-immunization-schedule-ages-6months-older.pdf">https://www.cdc.gov/vaccines/covid-19/downloads/COVID-19-immunization-schedule-ages-6months-older.pdf</a></li> </ul>
<b>Pre-vaccination counseling</b>	<p>Prior to vaccination:</p> <ul style="list-style-type: none"> <li>Provide the vaccine-specific Fact Sheet for Recipients and Caregivers Pfizer-BioNTech (<a href="https://www.fda.gov/media/144413/download">https://www.fda.gov/media/144413/download</a>), Moderna (<a href="https://www.fda.gov/media/144637/download">https://www.fda.gov/media/144637/download</a>), Janssen (<a href="https://www.fda.gov/media/146304/download">https://www.fda.gov/media/146304/download</a>), Novavax (<a href="http://www.novavaxcovidvaccine.com">www.novavaxcovidvaccine.com</a>)</li> <li>Screen for contraindications and precautions. CDC's Prevaccination Screening Form and Guidance document can be found at <a href="http://www.cdc.gov/vaccines/covid-19/info-by-product/index.html">www.cdc.gov/vaccines/covid-19/info-by-product/index.html</a>.</li> <li>Inform vaccine recipients mRNA or Novavax COVID-19 vaccines are recommended over Janssen COVID-19 Vaccine.</li> <li>Counsel vaccine recipients, parents, or guardians about expected reactions post-vaccination (e.g., pain and swelling at the injection site, fever, fatigue, headaches).</li> <li>Inform mRNA or Novavax vaccine recipients especially males ages 12-39 years, of the rare risk of myocarditis and/or pericarditis following receipt of these COVID-19 vaccines and the benefit of COVID-19 vaccination in reducing the risk of severe outcomes from COVID-19.<sup>†</sup> Counseling should also include the need to seek care if symptoms of myocarditis or pericarditis occur after vaccination, particularly in the week following vaccination. For more information see: <a href="https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us.html#myocarditis-pericarditis">https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us.html#myocarditis-pericarditis</a>.</li> <li>Janssen COVID-19 Vaccine is authorized for persons 18 years of age and older in certain limited situations due to safety considerations.</li> <li>Inform vaccine recipients interested in or receiving Janssen COVID-19 Vaccine of the risk and symptoms of thrombosis with thrombocytopenia syndrome (TTS), as well as the need to seek immediate medical care should symptoms develop after receiving Janssen vaccine. For more information see: <a href="https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us-appendix.html#appendix-a">https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us-appendix.html#appendix-a</a>.</li> </ul>
<b>Interchangeability of vaccines</b>	<ul style="list-style-type: none"> <li>In general, the same COVID-19 vaccine product (Pfizer-BioNTech, Moderna, Novavax) should be used for all doses in the primary series. In exceptional situations when the previous product cannot be determined/not available or if a person is unable to complete a series with the same COVID-19 vaccine (due to a contraindication), any age-appropriate mRNA COVID-19 vaccine may be used (administer at a minimum interval of 28 days).</li> <li>Any age-appropriate mRNA COVID-19 vaccine can be used for the booster dose(s); mRNA vaccines are recommended. Janssen COVID-19 Vaccine can be administered to persons 18 years of age and older in certain limited situations due to safety considerations and is not authorized for the 2nd booster dose. The Novavax COVID-19 Vaccine cannot be used for any booster dose. (<a href="https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us.html#timing-spacing-interchangeability">https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us.html#timing-spacing-interchangeability</a>).<sup>†</sup></li> </ul>
<b>Coadministration with other vaccines</b>	<ul style="list-style-type: none"> <li>COVID-19 vaccines may be administered on the same day as other vaccines.</li> <li>Administer each injection in a different injection site.</li> </ul>
<b>Contraindications</b>	<p>History of:</p> <ul style="list-style-type: none"> <li>Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a component of the COVID-19 vaccine</li> <li>A known diagnosed allergy to a component of the COVID-19 vaccine</li> <li>For the Janssen COVID-19 Vaccine, TTS following receipt of a previous Janssen COVID-19 Vaccine (or other COVID-19 vaccines not currently authorized or approved in the United States that are based on adenovirus vectors, e.g., AstraZeneca)<sup>‡</sup></li> </ul>

\* See Interim Clinical Considerations for Use of COVID-19 Vaccines Currently Approved or Authorized in the United States at: [www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us.html#recommendations](https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us.html#recommendations) for detailed guidance.

† Although CDC provides considerations for a mixed series in exceptional circumstances for a primary or additional dose, this is still considered an administration error that requires VAERS reporting. Heterologous booster doses are allowed (mRNA vaccines preferred) and are not considered a vaccine error.

‡ Additionally, people with a history of an episode of immune-mediated syndrome characterized by thrombosis and thrombocytopenia, such as spontaneous or classic HIT, should not receive Janssen COVID-19 Vaccine. These people should receive an mRNA COVID-19 vaccine.

# Summary Document for Interim Clinical Considerations

for Use of COVID-19 Vaccines Currently Authorized or Approved in the United States



## All currently authorized or approved COVID-19 vaccines

<b>Precautions</b>	<ul style="list-style-type: none"> <li>■ History of immediate allergic reaction to any other vaccine or injectable therapy (i.e., intramuscular, intravenous, or subcutaneous vaccines or therapies [excluding subcutaneous immunotherapy for allergies, i.e., “allergy shots”])</li> <li>■ History of multisystem inflammatory syndrome in children (MIS-C) or multisystem inflammatory syndrome in adults (MIS-A)</li> <li>■ History of an immediate (within 4 hours of exposure) non-severe allergic reaction after a dose of one type of COVID-19 vaccine have a precaution to the same type of COVID-19 vaccine</li> <li>■ Allergy-related contraindication to one type of COVID-19 vaccine have a contraindication or precaution to the other types of COVID-19 vaccines.</li> <li>■ Moderate or severe acute illness, with or without fever</li> <li>■ History of myocarditis or pericarditis after a dose of an mRNA or Novavax COVID-19 vaccine</li> <li>■ For Janssen COVID-19 Vaccine, a history of Guillain-Barré syndrome<sup>5</sup></li> </ul>
--------------------	---

## Considerations for all FDA-authorized or -approved COVID-19 vaccines

<b>Persons receiving HCT and CAR-T-cell therapy</b>	<ul style="list-style-type: none"> <li>■ If received doses of COVID-19 vaccine prior to or during HCT or CAR-T cell therapy, should be revaccinated for any doses received before or during treatment at least 3 months (12 weeks) after transplant or CAR-T-cell therapy</li> </ul>
<b>Persons who are moderately or severely immunocompromised</b>	<ul style="list-style-type: none"> <li>■ In most cases can receive a COVID-19 vaccine. Additional doses are recommended for this population for some vaccines.</li> <li>■ See the Interim COVID-19 Immunization Schedule for Ages 6 Months or Older at <a href="https://www.cdc.gov/vaccines/covid-19/downloads/COVID-19-immunization-schedule-ages-6months-older.pdf">https://www.cdc.gov/vaccines/covid-19/downloads/COVID-19-immunization-schedule-ages-6months-older.pdf</a></li> </ul>
<b>Persons receiving immunosuppressive therapies</b>	<ul style="list-style-type: none"> <li>■ Whenever possible, COVID-19 vaccines should be administered at least 2 weeks before initiation or resumption of immunosuppressive therapies</li> </ul>
<b>SARS-CoV-2 infection</b> <ul style="list-style-type: none"> <li>■ Current infection</li> <li>■ History of previous infection</li> <li>■ Exposed to an infected person</li> </ul>	<p>COVID-19 vaccination is recommended for everyone ages 6 months and older, regardless of a history of symptomatic or asymptomatic SARS-CoV-2 infection.</p> <ul style="list-style-type: none"> <li>■ Defer vaccination until person has recovered from acute illness and criteria have been met for them to discontinue isolation.</li> <li>■ People who recently had SARS-CoV-2 infection may consider delaying their next COVID-19 dose by 3 months from symptom onset or positive test (if infection was asymptomatic).</li> <li>■ Viral testing to assess for acute SARS-CoV-2 infection or serologic testing to assess for prior infection is not recommended for the purpose of vaccine decision-making.</li> <li>■ Additional information at: <a href="https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us.html#infection">https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us.html#infection</a></li> <li>■ COVID-19 vaccination is not recommended for post-exposure prophylaxis.</li> </ul>
<b>Persons with history of multisystem inflammatory syndrome (MIS-C and MIS-A) from SARS-CoV-2 infection</b>	<ul style="list-style-type: none"> <li>■ COVID-19 vaccines can be given; for children and adolescents wait at least 90 days after an MIS-C diagnosis.</li> <li>■ For persons who have had MIS-C or MIS-A from SARS-CoV-2 infection who have not yet received COVID-19 vaccine or who developed MIS-C or MIS-A after COVID-19 vaccination, a conversation between the vaccine recipient, guardian, and clinical team or specialist to discuss benefits and risks of receiving a COVID-19 vaccine is encouraged.</li> <li>■ Clinical recovery, including return to normal cardiac function, is an important factor when considering COVID-19 vaccination. Additional information at: <a href="https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us.html#COVID19-vaccination-misc-misa">https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us.html#COVID19-vaccination-misc-misa</a></li> </ul>

<sup>5</sup> People who develop GBS within 6 weeks after receipt of Janssen COVID-19 Vaccine should not receive another dose of Janssen COVID-19 Vaccine. These people should receive an mRNA COVID-19 vaccine.

# Summary Document for Interim Clinical Considerations

for Use of COVID-19 Vaccines Currently Authorized or Approved in the United States



## Considerations for all FDA-authorized or -approved COVID-19 vaccines

**Persons who received passive antibody therapy (convalescent plasma/ monoclonal antibodies)**

- COVID-19 vaccination can be given at any interval following receipt of passive antibody therapy.
- Persons should wait 2 weeks after COVID-19 vaccination before receiving tixagevimab/cilgavimab (EVUSHELD) for pre-exposure prophylaxis.

**Persons who are pregnant, breastfeeding, trying to get pregnant, or might become pregnant in the future**

- Are recommended to receive a COVID-19 vaccine primary series, additional mRNA doses (if indicated), and a booster dose(s).

## Considerations for mRNA vaccines and Novavax

**Persons with a history of myocarditis or pericarditis**

- Development of myocarditis or pericarditis after a dose of an mRNA or Novavax COVID-19 vaccine is a precaution to a subsequent dose of any COVID-19 vaccine.
- If after a risk assessment the decision is made to receive a subsequent COVID-19 vaccine dose, the person should wait until after their episode has resolved.
- For information on potential use of Janssen COVID-19 Vaccine in this situation, see Appendix A at [www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us-appendix.html#appendix-a](https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us-appendix.html#appendix-a)
- Persons who have a history of myocarditis or pericarditis unrelated to mRNA or Novavax COVID-19 vaccination may receive any age-appropriate COVID-19 vaccine after the episode of myocarditis or pericarditis has resolved.
- For more information see: <https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us.html#myocarditis-pericarditis>

## Considerations for Janssen COVID-19 Vaccine

Janssen COVID-19 Vaccine is authorized for adults ages 18 years and older in certain limited situations due to safety considerations. For more information, see <https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us-appendix.html#appendix-a>

**Persons with a history of Guillain-Barré syndrome (GBS)**

- A history of GBS is a precaution for receipt of Janssen COVID-19 Vaccine. An mRNA or Novavax vaccine is recommended.
- Persons who develop GBS within 6 weeks of Janssen COVID-19 vaccination should only receive an mRNA COVID-19 vaccine for subsequent doses.

**Persons with a history of thrombosis with thrombocytopenia syndrome (TTS)**

- It is contraindicated to administer Janssen COVID-19 Vaccine to persons with a history of TTS following receipt of the Janssen COVID-19 Vaccine or any other adenovirus vector-based COVID-19 vaccines (e.g., AstraZeneca's COVID-19 Vaccine).
- These persons should receive a dose of an mRNA COVID-19 vaccine as a booster dose at least 2 months (8 weeks) following their dose of the Janssen COVID-19 Vaccine and after their clinical condition has stabilized.

**Persons with a history of heparin-induced thrombocytopenia (HIT)**

- Persons with a history of an episode of an immune-mediated syndrome characterized by TTS, such as a spontaneous or classic HIT, should not receive Janssen COVID-19 Vaccine.
- These persons should receive an mRNA or Novavax COVID-19 vaccine.

# Summary Document for Interim Clinical Considerations

for Use of COVID-19 Vaccines Currently Authorized or Approved in the United States



## General COVID-19 Vaccination Information

<p><b>Persons vaccinated outside the United States</b></p>	<ul style="list-style-type: none"> <li>■ The recommendations for people vaccinated outside the United States depend on the number and type of vaccine(s) received for the primary series and booster doses, whether the primary series was completed, and whether a booster dose was received. Current guidance can be found at: <a href="https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us.html#appendix-a">https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us.html#appendix-a</a></li> </ul>
<p><b>Post-vaccination observation periods</b></p>	<p><b>30 minutes</b> – people with a history of:</p> <ul style="list-style-type: none"> <li>■ A contraindication to different type of COVID-19 vaccine product (i.e., mRNA or viral vector COVID-19 vaccines)</li> <li>■ Immediate (within 4 hours of exposure) non-severe allergic reaction to a COVID-19 vaccine</li> <li>■ Immediate allergic reaction of any severity to non-COVID-19 vaccine(s) or injectable therapies</li> <li>■ Anaphylaxis due to any cause</li> </ul> <p><b>15 minutes</b> – all other persons</p>
<p><b>SARS-CoV-2 antibody testing</b></p>	<ul style="list-style-type: none"> <li>■ Antibody testing is not recommended for vaccine decision-making or to assess immunity following vaccination.</li> </ul>
<p><b>Reporting requirements</b></p>	<p>Adverse events that occur following COVID-19 vaccination should be reported to VAERS (<a href="https://vaers.hhs.gov/">https://vaers.hhs.gov/</a>). COVID-19 providers are required to report:</p> <ul style="list-style-type: none"> <li>■ Vaccine administration errors</li> <li>■ Serious adverse events</li> <li>■ Cases of Multisystem Inflammatory Syndrome</li> <li>■ Cases of COVID-19 that result in hospitalization or death</li> </ul>